



VP INNOVATO HOLDINGS

iCervico

A disruptive new concept in implantology




Table of Contents

iCervico system

Introduction of the iCervico main components	3
--	---

iCervico healers

Shapes and sizes	4
------------------	---

Rotation and customization	5
----------------------------	---

Prosthetic components

Introduction of the i-base	6
----------------------------	---

iCervico sleeves & accessories	7
--------------------------------	---

Clinical protocols

• Healing stage	8
- Utilizing prefabricated healers	9
- Utilizing the Cervico Mold	10
• Single unit prosthesis	11
• Multiple unit prosthesis	15

iCervico components

AR / R - Sleeve



Retention screw



iCervico Healer



i-base




iCervico Healers

		Anteriors	Premolars	Lower Molars	Upper molars
Large	Off-set				
	Centered				
Medium	Off-set				
	Centered				
Small	Off-set				
	Centered	 6 mm	 7 mm		

iCervico Healer

Free rotation and customization of the iCervico healer.

The iCervico healer can be oriented ideally on the i-base independent of the implant prosthetic connection / interface orientation, without affecting the digital protocol.



i-base

4 scan mark surfaces

Direct scanning in surgical and / or prosthetic phase, eliminating the need for scan posts

Up to 3x more surface area than an external hex implant

Reduced stress on prosthesis retention screw

Quad-surface geometry with 10° bevel

Superior fit of prosthesis, eliminating the marginal micro-gap and prosthesis mobility

Variety of cuff heights

Adaptability to implant depth placement and supra-crestal soft tissue height.

- 1.5mm
- 2.5mm
- 3.5mm

Concave design shoulder with 15° emergence angle

Favorable soft and hard tissue response

One-piece

Absence of chimney eliminates risk of bacteria infiltration to implant platform

Regulating 30° angulation of implant & an additional 30° angulation of the screw access bore in 360° arch



Why i-base

One digital abutment for all stages of treatment

- Direct scan post
- Superior design for aesthetics
- Innovative design for prosthesis retention
- Suitable for both single and multiple prosthesis
- Functional with both analogic and digital protocols

Abutment level prosthetic protocol

- No unnecessary transgingival component disconnections that can lead to tissue recession
- Easy evaluation of prosthetic component fit during all restorative stages; simplifying the restorative process of conical connection implants.
- Allows up to 30° angulation correction (short AR/R- sleeve)

Multi purpose sleeves

- **Titanium sleeve prosthesis interface** - ensuring ideal prosthesis fit and retention
- **Scan-post extension** - for cases where the geometry of the i-base cone is not easily scannable


AR - Sleeve | Anti - Rotational

Suitable for single unit prosthesis



Quad-surface conical geometry

Allows precise and anti-rotational fit onto the i-base



Two external flat surfaces for:

- Stable and precise fit into the prosthesis
- Scan mark surface


R - Sleeve | Rotational

Suitable for multiple unit prosthesis



Circular geometry

Provides ease for multiple unit prosthesis path of insertion



Embedded cut-mark

allows for quick and accurate alteration of sleeve height.



Retention Screw

Innovative retaining screw made of Ti6Al4V alloy with a diameter of M1.6 (similar diameter with TI-BASE retaining screw) allowing a screw torque of 30 N/cm.

Corrects up to 30 degrees of angle in a 360 degree direction.

ACCESSORIES | Lab analog - Impression post - ISQ

iCervico lab analog



- One analog for all i-base options
- Compatible with both analogic and digital protocols
- Innovative design for accurate positioning into the 3D printed models

Impression post



- Suitable for both single & multiple unit cases
- Suitable for both open and closed tray impression techniques.
- Scannable version available

Smart peg - ISQ



- iCervico Smart Peg for ISQ measurements.
- Easily evaluate implant stability without disturbing the soft tissue seal.

Clinical protocol



Healing stage



VP INNOVATO HOLDINGS

Prefabricated iCervico healers

Clinical protocol

Utilize the desired iCervico healer directly on top of the i-base.




Chairside iCervico healers

Clinical protocol

Make chair-side your own iCervico healers with the use of CERVICO MOLD by customising the R-sleeves with either of the following workflows:


- Use the R-sleeve to create an anatomical abutment which can be then customized into a temporary prosthesis
- Cut the R-sleeve extension and fabricate an anatomical healing abutment



iCervico healing abutment

iCervico temporary abutment

Clinical protocol




Single unit prosthesis

Clinical Protocol

Single unit prosthesis




3 different Scanning / Impression stage options



Clinical Protocol | Single unit prosthesis




Register opposite arch and occlusion, and share with the Lab.




Reinstall the iCervico healer after scan / impression

Lab Protocol



Utilize the iCervico digital library and declare the single digital scan as:

1. Jaw
2. Soft Tissue
3. Scan Post Proceed



Chose on the library the type of prosthesis:

a. Single unit

- AR-sleeve, long
- AR-sleeve, short

b. Bridge


- R-sleeve, long
- R-sleeve, short



Proceed with prosthesis design

Clinical Protocol | Single unit prosthesis

Lab stage



Adjust screw
channel angulation
if necessary
(up to 30°) with
short AR-sleeve



Utilize the
appropriate
height
AR-sleeve




Fabricate the
provisional
or final prosthesis




Cement the
AR-sleeve
into the
prosthesis
extraorally


Delivery stage



Remove the
iCervico healer




Install the screw-retained
prosthesis
on the i-base



Apply 30 N/cm torque


Clinical protocol





Multiple unit prosthesis

Clinical Protocol


Multiple unit prosthesis




After completion of osseo & muco-integration
remove the iCervico healer in order to proceed
with digital or analogic impression stage




3 different Scanning / Impression stage options



a. Direct scan of i-bases




b. Install AR-sleeves and scan





c. Install Impression posts coupled with the impression screw and take analogic impression

*The use of AR-sleeves provides the anti-rotational position of the i-base while the use of R-sleeves does not.

Reinstall iCervico healers until prosthesis delivery



Lab stage



Utilize the iCervico digital library and declare the single digital scan as:

1. Jaw
2. Soft Tissue
3. Scan Post

Chose on the library the type of prosthesis:

a. Single Unit

- AR-sleeve, long
- AR-sleeve, short

b. Bridge

- R-sleeve, long
- R-sleeve, short



Proceed with
prosthesis design

Clinical Protocol | Multiple unit prosthesis

Lab stage



(Optional)

Adjust screw channel angulation if necessary (up to 30°) with short R-sleeve



Utilize the appropriate height R-sleeves




Fabricate the provisional or final prosthesis




Cement the R-sleeves into the prosthesis extraorally


Delivery stage




Remove the iCervico healers




Install the screw-retained prosthesis on the i-bases and apply 30 N/cm torque



Rectangular



Ovoid



Triangular

Transition smoothly from your cervico profiler to the final prosthesis maintaining a biologically sound cervical zone



ariston dental

better answers

ATHENS

467 Mesogeion Ave., Bld. A,
153 43, T: +30 210 80 30 341
info@aristondental.com

W: aristondental.com

THESSALONIKI

13 Navarchou Kountouriotou Str.,
546 25, T: +30 2310 268 020
thessaloniki@aristondental.com

fb: ariston dental

LIMASSOL

75 Leontiou Maxaira Str.,
Office 203, 3025, Limassol,
T: +357 25322895
info@aristondental.com